

Effect of Time and Frequency Manipulation on Syllable Perception in Developmental Dyslexics

Ken I. McAnally*

Peter C. Hansen

Piers L. Cornelissen**

John F. Stein

University Laboratory of Physiology
Oxford University, England

Many people with developmental dyslexia have difficulty perceiving stop consonant contrasts as effectively as other people and it has been suggested that this may be due to perceptual limitations of a temporal nature. Accordingly, we predicted that perception of such stimuli by listeners with dyslexia might be improved by stretching them in time—equivalent to speaking slowly. Conversely, their perception of the same stimuli ought to be made even worse by compressing them in time—equivalent to speaking quickly. We tested 15 children with dyslexia on their ability to identify correctly consonant-vowel-consonant (CVC) stimuli that had been stretched or compressed in the time domain. We also tested their perception of the same CVC stimuli after the formant transitions had been stretched or compressed in the frequency domain. Contrary to our predictions, we failed to find any systematic improvement in their performance with either manipulation. We conclude that simple manipulations in the time and frequency domains are unlikely to benefit the ability of people with dyslexia to discriminate between CVCs containing stop consonants.

KEY WORDS: dyslexia, speech perception, auditory perception, reading disabilities, disorders

Many children fail to acquire competent reading skills despite adequate educational opportunity. A surprising number of these poor readers have normal or above normal abilities in other areas, so their reading problems are unexpected. Rutter and Yule (1975) attempted to capture this sense of unexpected reading failure by using statistical criteria. They defined children as specifically retarded readers (i.e., developmental dyslexics) if their reading ability was significantly lower than that predicted on the basis of age and IQ. Stanovich (1991) has criticized IQ discrepancy measures of developmental dyslexia because poor reading ability tends to be correlated with poor performance on IQ tests. Consequently, some authors have suggested that a mismatch between reading and spoken comprehension might provide a better measure (Gough & Tunmer, 1986). Nevertheless, the majority of published research has used some version of the

*Currently affiliated with the Air Operations Division, Aeronautical and Maritime Research Laboratory, Melbourne, Australia.

**Currently affiliated with the Psychology Department, Newcastle University, Newcastle upon Tyne, England.

IQ discrepancy measure to define developmental dyslexia. For the sake of consistency, therefore, we have used the same definition.

Many people with developmental dyslexia have difficulty perceiving consonant contrasts. Both single case and group studies have shown that people with dyslexia confuse the identity of a range of fricatives in spoken consonant-vowel (CV) syllables (Cornelissen, Hansen, Bradley, & Stein, 1996; Masterson, Hazan, & Wijayatilake, 1995). Furthermore, at least six studies have been published in which categorical perception of speech sounds was investigated in dyslexic individuals by using synthetic CV continua (/ba/-/da/ and /da/-/ga/, which differ in the frequency of the second formant transition; /bath/-/path/ and /sa/-/sta/, which differ on voice onset time) (Brandt & Rosen, 1980; Godfrey, Syrdal-Lasky, Millay, & Knox, 1981; Manis, McBride, Seidenberg, Doi, & Custodio, 1993; Reed, 1989; Steffens, Eilers, Gross-Glenn, & Jallad, 1992; Werker & Tees, 1987). Although people with dyslexia demonstrate categorical perception of synthetic CV continua containing stop consonants, the slopes of the boundary curves for the listeners with dyslexia tend to be less steep than normal. This suggests that the categorical boundaries between different speech sounds may be blurred in people with dyslexia. Thus, they may confuse phonetically similar speech sounds more readily than normal.

It has been proposed that people with dyslexia have deficits in temporal processing of acoustic information (Tallal, 1980) that may limit the accuracy of their identification of consonant-vowel F2 and F3 transitions (Tallal & Stark, 1981). In support of temporal processing deficits in dyslexia, Watson (1992) showed that reading-disabled college students were impaired in their ability to detect differences in the duration of tones but not in their ability to detect changes in frequency. If people with dyslexia have temporal processing deficits, their perception of speech stimuli might be improved by stretching these stimuli in time. Conversely, their perception of the same stimuli might be made even worse by compressing them in time. We examined the ability of dyslexic and nondyslexic children to identify synthetic stop consonant CVC stimuli that were compressed or

expanded in time, while retaining the original frequency content.

Watkins, Baldeweg, Richardson, and Gruzelier (1995) and McAnally & Stein (1996) have shown that adult dyslexics are more impaired in their discrimination of frequency than of either tone or gap duration. Therefore, we also investigated identification of synthetic CVC stimuli in which the range of the formant frequency transitions was compressed or expanded.

Method

Participants

Fifteen dyslexic boys who had been assessed by the same educational psychologist participated in the experiment (mean age was 182 months, standard deviation [*SD*] 11.8; mean IQ [British Ability Scales, BAS] was 108, *SD* 8.1). Diagnosis was based on standard IQ discrepancy criteria where reading age is two or more standard deviations below the mean for children of similar age and IQ. Fifteen boys from the same school with no history of reading problems also participated (mean age was 183 months, *SD* 10.8; mean IQ [BAS] 124, *SD* 11.3). All participants had normal hearing thresholds to pure tones between 125 Hz and 8 kHz. Independent of the psychologist's assessments, we administered two reading tests. Subjects were given a list of real words (Schonell, 1950) or nonsense words (Castles & Coltheart, 1993) and were instructed to read the lists as quickly and as accurately as possible. The participants with dyslexia made more errors and took longer to read both the word and nonword lists than did participants with no history of reading problems (Figure 1 and Table 1). In view of the group difference in IQ, we carried out an analysis of covariance on the reading error data taking both age and IQ into account. The children with dyslexia were significantly poorer readers than those without, even when age and IQ were taken into account (Table 2). It is clear from these data that the children with dyslexia had persistent reading problems.

The ease of identification of the baseline and manipulated synthetic speech stimuli was assessed using four adults who were ignorant of the aims of this study.

Table 1. Participant characteristics.

| | Dyslexic (<i>n</i> = 15) | | Nondyslexic (<i>n</i> = 15) | |
|----------------------------|---------------------------|---------------|------------------------------|---------------|
| | Mean | (<i>SD</i>) | Mean | (<i>SD</i>) |
| Word list: errors | 22.3 | (7.8) | 6.8 | (5.9) |
| Word list: time (s) | 227.6 | (181.8) | 100.0 | (25.4) |
| Nonword list: errors | 9.5 | (6.9) | 2.7 | (3.4) |
| Nonword list: time (s) | 55.2 | (20.4) | 28.4 | (7.1) |
| Chronological age (months) | 182.4 | (11.8) | 182.6 | (10.8) |
| IQ (BAS) | 107.8 | (8.1) | 124.4 | (11.3) |

Experimental Stimuli

We used a speech synthesizer (Klatt, 1980) at a sample rate of 10 kHz to generate a baseline set of consonant-vowel-consonant (CVC) stimuli. Synthesizer parameters were generated by the IPOX interface (Dirksen & Coleman, 1995) to synthesize an adult male voice. The stimuli contained all the stop consonants in either onset or rime positions separated by the vowel sound /b/

Figure 1. Two plots of the total number of reading errors that each subject made against the time it took them to complete the reading lists. The first plot is for the real word reading test and the second plot for the nonword reading test. Participants with dyslexia are represented by solid triangles while nondyslexic participants are represented by open circles.




Table 2. Analysis of covariance for word and nonword reading errors.

| | Explanatory variable | F ratio (df) | p value |
|----------------|----------------------|--------------|--------------|
| Word errors | Participant group | 22.1 (1,26) | $p < 0.0005$ |
| | Chronological age | 3.4 (1,26) | $p > 0.05$ |
| | IQ (BAS) | 0.1 (1,26) | $p > 0.5$ |
| Nonword errors | Participant group | 7.8 (1,26) | $p < 0.05$ |
| | Chronological age | 6.7 (1,26) | $p < 0.05$ |
| | IQ (BAS) | 0.3 (1,26) | $p > 0.5$ |

(/bɒk/, /pɒk/, /dɒk/, /tɒk/, /gɒk/, /kɒk/, /kɒg/, /kɒb/, /kɒp/, /kɒd/ and /kɒt/). An example of the synthesizer parameters for the stimulus /dɒk/ is given in Appendix A. Differences from the default output from IPOX are listed in Appendix B for /kɒb/ and /kɒp/. We applied a uniform stretch or compression to the time-course of all synthesizer parameters, allowing us to generate a time-stretched (x1.6) and a time-compressed (x0.6) version of each stimulus. This algorithm did not alter the frequencies of the fundamental or the formants but altered their rate of change. Informal listening showed the effect of this algorithm; it made the CVC syllables sound like someone

speaking either very quickly (time compression) or very slowly (time stretching).

We also generated stimuli in which the range of the transitions of formant frequency was either stretched or compressed, but the steady-state formant frequency of the vowel remained unchanged. However, their duration remained unchanged. We applied a frequency stretch or compression algorithm (around the steady-state value) to the formant frequencies of the same baseline set of synthesizer parameters to generate a frequency-stretched (x1.4) and a frequency-compressed (x0.5) version of each baseline stimulus. Informal listening showed that the effect of this algorithm was similar to the difference between someone enunciating clearly (frequency stretch) or lazily (frequency compression). Spectral plots of the four kinds of manipulated stimuli are shown in Figure 2.


Synthetic speech stimuli were played by an IBM compatible PC equipped with a 16-bit sound card (Sound Blaster 16, Creative Labs.) and presented diotically through headphones (Sennheiser HD520II) at a sound pressure of 70dB (with respect to 20 μ Pa).

Data Collection

Each trial was composed of a single interval in which a stimulus was chosen randomly (without replacement) from the set of stimuli. Listener responses were recorded using a multiple choice window presented on a computer monitor. The possible responses were represented on screen in a vertical array of 11 boxes, each of which contained one of the letter strings (bok, pok, dok, tok, gok, kok, kog, kob, kop, kod, and kot). Once participants had decided which of the stimuli they had heard, they selected the appropriate response box with the mouse. The computer logged the stimulus and the response. The positions of the letter strings in the array of boxes were randomized on every trial to control for any tendency for participants to prefer selecting particular box locations.

Before the experiment, participants were asked to read aloud the letter strings within each response box and occasionally they made mistakes. Participants who made mistakes were asked to repeat the task until they could read through the letter strings perfectly.

Figure 2. A: Frequency expanded version of /bɒk/. B: Frequency compressed version of /bɒk/. C: Time compressed version of /bɒk/. D: Time expanded version of /bɒk/. Darker shading represents increasing amplitude.



The experiment was run in 12 blocks under computer control (Canadian Speech Research Environment). The first block was a training condition during which subjects were familiarized with the synthetic speech stimuli. Participants listened to five repetitions of each of the 11 baseline CVC stimuli, presented in random order. Visual feedback was provided about whether the responses were correct. The next 10 blocks of trials provided the main experimental data. Each block comprised a presentation of each of the five versions of each CVC (baseline, time stretched, time compressed, frequency stretched and frequency compressed) in random order without replacement in a background of white noise (signal-to-noise ratio of 15 dB). Low-level noise was used to reduce the children's performance enough for us to measure any potential benefit from the time or frequency manipulations. If we had not added white noise, many children's performances might have been close to ceiling even with the baseline stimuli.

To control for possible difficulties that the children with dyslexia may have had reading the letter strings in each response box, the stimuli were presented visually as letter strings in the final block. Each of the 11 letter strings was presented 10 times. On each trial, the stimulus letter string appeared on screen for 500 ms before the response boxes were presented.

Results

Table 3 shows the proportion of correct responses in the visual control task, the noise free training condition, and the experimental condition for the baseline stimuli only. The high scores on the visual control task show that dyslexic and nondyslexic children were able to at least recognize the visual stimuli. The fact that performance was so good on the visual control task and that the children were able to read the visual stimuli aloud suggest that participants' responses were unlikely to be confounded by the problem of reading from the multiple choice response box. For the auditory stimuli presented in quiet, the adults scored 94.1% correct, showing the baseline stimuli to be well discriminable and identified. Chance performance in the task would have

Table 3. Percent correct scores on the speech tasks (baseline stimuli only) and visual control task.

| | Children | | Adults | |
|-------------|----------------|-----------------------------------|--------------------------------------|-----------------------------------|
| | Visual control | Spoken CVC in quiet with feedback | Spoken CVC in noise without feedback | Spoken CVC in quiet with feedback |
| Dyslexic | 91.8% | 70.4% | 54.9% | N/A |
| Nondyslexic | 96.1% | 76.5% | 63.9% | 94.1% |

resulted in 9% correct identification. The time and frequency modified syllables in noise were all also well identified by the adult listeners (baseline 84.1%, time compressed 63.6%, time stretched 69.1%, frequency compressed 75.2% and frequency stretched 73.6%), suggesting that these syllables did not cross perceptual boundaries following modification. Both dyslexic and nondyslexic children scored less well than the adults for the auditory stimuli presented in quiet and, as expected, the performance of both groups of children was impaired by the combination of white noise masking and the removal of feedback.


Table 1 showed that the mean IQ of the children with dyslexia was lower than the mean IQ of the controls. However, Table 4 shows that the correlation between IQ and the performance in the speech task (for the five kinds of stimuli) was always weak and insignificant. Therefore, IQ was excluded from the following analyses.

As a first step toward analyzing the effects of our interventions, we carried out separate repeated measures analyses of variance (ANOVA) of responses to the time and frequency modified stimuli, comparing responses of participants with or without dyslexia. Figure 3 shows a plot of the means and standard deviations for the ANOVAs. It is clear from these plots that neither expansion nor compression of time or frequency resulted in an improvement over baseline in either group of participants. The appropriate probability distribution for proportionate data is binomial because we were calculating the proportion of errors made. We therefore applied an arcsine transform (Snedecor & Cochran, 1967) to the data. For the time-manipulated stimuli, although the main effect of stimulus-type was significant ($F_{2,56} = 32.6, p < 0.0005$), neither the main effect of group (i.e., dyslexic vs. nondyslexic) nor the interaction between group and stimulus-type was significant ($F_{1,28} = 2.9, p > 0.1$, and $F_{2,56} = 0.56, p > 0.5$, respectively). For the frequency-manipulated stimuli we found similar results. Although the main effect of stimulus-type was significant ($F_{2,56} = 64.7, p < 0.0005$), neither the main effect of group (i.e., dyslexic vs. nondyslexic) nor the interaction between group and stimulus-type was significant ($F_{1,28} = 2.7, p > 0.1$, and $F_{2,56} = 0.64, p > 0.5$, respectively). Both time and frequency manipulation affected

Table 4. Correlations between IQ and the percentage correct responses in the speech task.

| Stimulus type | Correlation coefficient | p value |
|----------------------|-------------------------|------------|
| Baseline | 0.36 | $p > 0.05$ |
| Frequency compressed | 0.29 | $p > 0.1$ |
| Frequency stretched | 0.31 | $p > 0.1$ |
| Time compressed | 0.41 | $p > 0.5$ |
| Time stretched | 0.28 | $p > 1.1$ |

Figure 3. Plots of the percentage correct responses comparing children with and without dyslexia when listening to the time and frequency compressed or expanded stimuli, embedded in white noise. Error bars represent 1 *SD*. The x-axis represents the expansion factor so a value of 1.0 is equivalent to the baseline stimulus.



CVC identification, but contrary to our hypothesis they reduced performance. Moreover, these effects were the same whether children had dyslexia or not.



Our baseline stimuli varied over two dimensions: place of articulation and voice onset time (VOT). Similarly, in the studies of the categorical perception of CV sounds in people with dyslexia (see earlier), both place of articulation and VOT were varied. These studies showed that people with dyslexia exhibited categorical boundaries that were less sharp than normal. Hence, in our study, it is possible that time stretching might have improved the perception of isolated pairs of CVCs that were differentiated by a single phonetic feature like VOT (e.g., /bɒk/ vs. /pɒk/). Yet such an effect could have been masked in the above analysis by being averaged together with the more common effect of degraded performance for the majority of CVCs. Therefore, we sought isolated effects of this kind by looking at the consequences of our manipulations on the identification of the individual phonetic features: place of articulation and VOT. For place of articulation, we counted the percentage of correct responses within six phonetic categories: labial onsets (/bɒk/ and /pɒk/), alveolar onsets (/dɒk/ and /tɒk/), palatal onsets (/gɒk/ and /kɒk/), labial rimes (/kɒb/ and /kɒp/), alveolar rimes (/kɒd/ and /kɒt/), and palatal rimes (/kɒg/ and /kɒk/). For VOT, we counted the percentage of correct responses within four phonetic categories: voiced onsets (/bɒk/, /dɒk/, and /gɒk/), unvoiced onsets (/kɒk/, /pɒk/, and /tɒk/), voiced rimes (/kɒb/, /kɒd/, and /kɒg/), and unvoiced rimes (/kɒp/, /kɒt/, and /kɒt/). Within each category, we plotted the percentage of correct responses as a function of stimulus compression or expansion as shown in Figures 4, 5, 6, and 7. In each plot, a value on the x-axis of one represents the baseline stimuli, values less than one represent compression, and values greater than one represent expansion. We looked for improvement in performance over the baseline as a consequence of time or frequency manipulation. As can be seen in Figures 4, 5, 6, and 7, we could find none for either dyslexic or nondyslexic listeners. Instead, both time and frequency stretching tended to have a detrimental effect on identification of these CVCs.

Finally, we analyzed the data individual by individual to try to find any children whose perception of CVCs may have benefited from time or frequency manipulation. We selected cases where time or frequency stretching produced an increase in the proportion of correct responses of 25% or more. Tables 5 and 6 show the instances where such improvement was recorded for place of articulation. For VOT, there was only one case of improved identification: The identification of unvoiced onsets was improved in one individual by time compression. To our surprise, compression as well as expansion in both the time and frequency domains produced examples of improvement in individuals from both listener groups. However, there did not seem to be any particular condition that was selectively beneficial to children with dyslexia.



Figure 4. Plots of the percentage correct responses of the children with dyslexia when listening to the time and frequency compressed or expanded stimuli, embedded in white noise. The data are plotted separately for the phonetic features: labial onsets (/bɒk/ and /pɒk/), alveolar onsets (/dɒk/ and /tɒk/), palatal onsets (/gɒk/ and /kɒk/), labial rimes (/kɒb/ and /kɒp/), alveolar rimes (/kɒd/ and /kɒt/), and palatal rimes (/kɒg/ and /kɒk/). The x-axis represents the expansion factor so a value of 1.0 is equivalent to the baseline stimulus.

Dyslexics

Frequency compression/expansion



Time compression/expansion



Discussion

Based on the results from previous studies that show children with dyslexia to have difficulty in processing rapid acoustic stimuli (Tallal, 1980), we predicted that time expansion ought to improve the ability of such children to discriminate between phonetically confusable spoken syllables. However, the present results fail to support these predictions for CVC syllables containing stop consonants. Stollman, Kapteyn, and Sleswijk (1994) have measured the speech recognition threshold in noise (SRT) for children with language impairment listening to normal and time-manipulated speech. As expected, the SRT increased with time compression, but contrary to the prediction based on a temporal processing deficit in children with



language impairment (Tallal & Piercy, 1973), the SRT was not improved with time expansion. Stollman et al.'s findings are consistent with those of the present study of children with dyslexia.

None of our manipulations (time or frequency, expansion or compression) improved overall CVC identification in either dyslexic or nondyslexic children. Nevertheless, a more detailed analysis of our results showed that the perception of certain phonetic features was improved in some individuals not only by time compression or expansion, but also by frequency compression or expansion. Unfortunately, these results did not apply systematically to any one individual; although a particular manipulation may have improved the perception



Figure 5. Plots of the percentage correct responses of the nondyslexic children when listening to the time and frequency compressed or expanded stimuli, embedded in white noise. As in Figure 3, the data are plotted separately for the phonetic features: labial onsets (/bɒk/ and /pɒk/), alveolar onsets (/dɒk/ and /tɒk/), palatal onsets (/gɒk/ and /kɒk/), labial rimes (/kɒb/ and /kɒp/), alveolar rimes (/kɒd/ and /kɒt/), and palatal rimes (/kɒg/ and /kɒk/). The x-axis represents the expansion factor so a value of 1.0 is equivalent to the baseline stimulus.

Nondyslexics

Frequency compression/expansion



Time compression/expansion




of some phonetic features, the same manipulation degraded the perception of other features. Moreover, we found isolated examples of improved performance in individuals from both listener groups. Therefore, it seems that manipulating either the time or the frequency domain as we have done is unlikely to lead to a dramatic improvement in the perception of stop consonant CVC syllables by listeners with dyslexia.

There was a significant difference in IQ (BAS) between the participant groups in the present study. This factor was statistically taken into account to ensure that the children with dyslexia were indeed worse readers than the group of nondyslexic children. However, this difference was unlikely to have had any influence on the comparison between dyslexic and nondyslexic individuals on CVC identification because there was no correlation

Figure 6. Plots of the percentage correct responses of the children with dyslexia when listening to the time and frequency compressed or expanded stimuli, embedded in white noise. The data are plotted separately for the phonetic features: voiced onsets (/bɔk/, /dɔk/, and /gɔk/), unvoiced onsets (/kɔk/, /pɔk/, and /tɔk/), voiced rimes (/kɔb/, /kɔd/, and /kɔg/), and unvoiced rimes (/kɔk/, /kɔp/, and /kɔt/). The x-axis represents the expansion factor so a value of 1.0 is equivalent to the baseline stimulus.

Dyslexics

Frequency compression/expansion



Time compression/expansion





Figure 7. Plots of the percentage correct responses of the nondyslexic children when listening to the time and frequency compressed or expanded stimuli, embedded in white noise. As in Figure 6, the data are plotted separately for the phonetic features: voiced onsets (/bɔk/, /dɔk/, and /gɔk/), unvoiced onsets (/kɔk/, /pɔk/, and /tɔk/), voiced rimes (/kɔb/, /kɔd/, and /kɔg/), and unvoiced rimes (/kɔk/, /kɔp/, and /kɔt/). The x-axis represents the expansion factor so a value of 1.0 is equivalent to the baseline stimulus.

Nondyslexics

Frequency compression/expansion



Time compression/expansion




Table 5. Individual analysis of data from children with dyslexia where improvement is $\geq 25\%$.

| Mode | Phonetic feature | Compression | | Expansion | |
|-----------|------------------|---------------|-------|-----------|-------|
| | | Onsets | Rimes | Onsets | Rimes |
| Time | Labial | DXM | GHR | DXM, MJB | |
| | Alveolar | | EMW | | WJA |
| | Palatal | | | | |
| Frequency | Labial | | | DXM | |
| | Alveolar | DXM, GHR, MJB | TJT | | WJA |
| | Palatal | WRC | | | |

Table 6. Individual analysis of data from children without dyslexia where improvement is $\geq 25\%$.

| Mode | Phonetic feature | Compression | | Expansion | |
|-----------|------------------|-------------|-------|-----------|-------|
| | | Onsets | Rimes | Onsets | Rimes |
| Time | Labial | EWT,RMM,TCF | CMS | JMW,RMM | |
| | Alveolar | | | | |
| | Palatal | | | | |
| Frequency | Labial | EMB | | | EWT |
| | Alveolar | | | | |
| | Palatal | SRH | CMS | | |

between IQ and performance on the speech task.

Recently, Tallal and colleagues reported a significant improvement in the language comprehension abilities of children with language impairment who had been given extensive training with both adaptive temporal discrimination exercises as well as acoustically modified speech (Merzenich et al., 1996; Tallal et al., 1996). It is unclear which aspect of the training regime in their study (i.e., the temporal discrimination exercises or the modified speech) contributed to the improvement in the children's language ability. Furthermore, acoustic modification of the speech was a two-stage process. The signal was initially slowed by 50% and then amplitude modulations between 3Hz and 30Hz in the narrowband filtered signal were amplified. If long-term exposure to the acoustically modified speech had been responsible for improving speech perception, it is unclear which of these processing stages (i.e., time expansion or amplitude modulation) was the more potent. The results from the present study suggest that limited exposure of children with dyslexia to time-stretched synthetic CVC syllables did not improve their ability to identify the stimuli correctly. However, McAnally and Stein (1997) have shown that the scalp potential evoked by amplitude modulation of tones is significantly smaller in adults with dyslexia than in matched control listeners. Therefore, as with the amplitude manipulations of stimuli by Tallal et al. (1996), it is possible that the selective emphasis of narrowband AM may be more potent than slowing of the speech signal in improving speech perception in listeners with dyslexia.

Acknowledgments

The authors thank staff and students of Bloxham school for their participation and John Coleman of Oxford University Phonetics Department. This work was supported by the McDonnell-Pew Centre for Cognitive Neuroscience and the Rodin Remediation Academy.

References

- Brandt, J., & Rosen, J. J.** (1980). Auditory phonemic perception in dyslexia: Categorical identification and discrimination of stop consonants. *Brain and Language, 9*, 324-337.
- Castles, A., & Coltheart, M.** (1993). Varieties of developmental dyslexia. *Cognition, 47*, 149-180.
- Cornelissen, P. L., Hansen, P. C., Bradley, L., & Stein, J. F.** (1996). Analysis of perceptual confusion between 9 sets of consonant-vowel sounds in normal and dyslexic adults. *Cognition, 59*, 275-306.
- Dirksen, A., & Coleman J.** (1995). IPO/Oxford: All-prosodic Speech Synthesis Architecture. Software available from the Oxford University Phonetics Department.
- Godfrey, J. J., Syrdal-Lasky, A. K., Millay, K. K., & Knox, C. M.** (1981). Performance of dyslexic children on speech perception tests. *Journal of Experimental Child Psychology, 32*, 401-424.
- Gough, P. B., & Tunmer, W. E.** (1986). Decoding, reading and reading disability. *Remedial and Special Education, 7*, 6-10.
- Klatt, D. H.** (1980). Software for a cascade/parallel formant synthesiser. *Journal of the Acoustical Society of America, 67*, 971-995.

- Manis, F., McBride, C., Seidenberg, M., Doi, L., & Custodio, R.** (1993). *Speech perception and phonological awareness in normal and disabled readers*. Poster presentation at meeting of the SRCD, New Orleans, LA.
- Masterson, J., Hazan, V., & Wijayatilake, L.** (1995). Phonemic processing problems in developmental phonological dyslexia. *Cognitive Neuropsychology*, *12*, 233–259.
- McAnally, K. I., & Stein, J. F.** (1996). Auditory temporal processing in dyslexia. *Proceedings of the Royal Society of London Series B* *263*, 961–965.
- McAnally, K. I., & Stein, J. F.** (1997). Scalp potentials evoked by amplitude-modulated tones in dyslexia. *Journal of Speech, Language, and Hearing Research*, *40*, 939–945.
- Merzenich, M. M., Jenkins, W. M., Johnston, P., Schreiner, C., Miller, S. L., & Tallal, P.** (1996). Temporal processing deficits of language-learning impaired children ameliorated by training. *Science*, *271*, 77–81.
- Reed, M. A.** (1989). Speech perception and the discrimination of brief auditory cues in reading disabled children. *Journal of Experimental Child Psychology*, *48*, 270–292.
- Rutter, M., & Yule, W.** (1975). The concept of specific reading retardation. *Journal of Child Psychology and Psychiatry*, *16*, 181–197.
- Schonell, F.J.** (1950). *Diagnostic and attainment testing*. London: Oliver & Boyd.
- Snedecor, G. W., & Cochran, W. G.** (1967) *Statistical methods*. Ames, IA: The Iowa State University Press.
- Stanovich, K.E.** (1991). The theoretical and practical consequences of discrepancy definitions of dyslexia. In M. J. Snowling & M. E. Thomson (Eds.) *Dyslexia: Integrating theory and practice* (p. 327). London and New Jersey: Whurr Publishers Ltd.
- Steffens, M. L., Eilers, R. E., Gross-Glenn, K., & Jallad, B.** (1992). Speech perception in adult subjects with familial dyslexia. *Journal of Speech and Hearing Research*, *35*, 192–200.
- Stollman, M. H. P., Kapteyn, T. S., & Sleeswijk, B. W.** (1994). Effect of time-scale modification of speech on the speech recognition threshold in noise for hearing-impaired and language-impaired children. *Scandinavian Audiology*, *23*, 39–46.
- Tallal, P.** (1980). Auditory temporal perception, phonics, and reading disabilities in children. *Brain and Language*, *9*, 182–198.
- Tallal, P., & Piercy, M.** (1973). Deficits of non-verbal auditory perception in children with developmental aphasia. *Nature*, *241*, 468–469.
- Tallal, P., & Stark, R. E.** (1981). Speech acoustic-cue discrimination abilities of normally developing and language-impaired children. *Journal of the Acoustical Society of America*, *69*, 568–533.
- Tallal, P., Miller, S. L., Bedi, G., Byma, G., Wang, X. Q., Nagarajan, S. S., Schreiner, C., Jenkins, W. M., & Merzenich, M. M.** (1996). Language comprehension in language-learning impaired children improved with acoustically modified speech. *Science*, *271*, 81–84.
- Watkins, S., Baldeweg, T., Richardson, A. J., & Gruzelier, J.** (1995). *Auditory mismatch evoked potentials in dyslexia: Evidence for a dysfunction in automatic feature detection for pitch*. Twenty-third annual meeting of the British Psychophysiology Society, Keele, U.K.
- Watson, B. U.** (1992). Auditory temporal acuity in normally achieving and learning-disabled college students. *Journal of Speech and Hearing Research*, *35*, 148–156.
- Werker, J. F., & Tees, R. C.** (1987). Speech perception in severely disabled and average reading children. *Canadian Journal of Psychology*, *41*(1), 48–61.

Received August 29, 1996

Accepted February 17, 1997

Contact author: Piers Cornelissen, Psychology Department, Ridley Building, Newcastle University, Newcastle upon Tyne, NE1 7RU, U.K.

Appendix A. Klatt parameter values for the baseline stimulus /dɒk/. The first number given is the parameter value and the second number, if present, is time (ms). Parameter labels are as in Klatt (1980).

| | | | | | | | | | |
|--------|--------|--------|-------|-------|-----|------|-----|------|------|
| SW | FGP | BGP | FGZ | BGZ | B4 | F5 | B5 | F6 | B6 |
| 0 | 0 | 100 | 1500 | 6000 | 250 | 3750 | 200 | 4900 | 1000 |
| FNP | BNP | BNZ | BGS | SR | NWS | G0 | NFC | AC | |
| 250 | 100 | 100 | 200 | 10000 | 50 | 48 | 5 | 0 | |
| AV | AF | AH | AVS | | | | | | |
| 0,5 | 0,5 | 0,5 | 0,5 | | | | | | |
| 0,300 | 0,275 | 0,560 | 0,830 | | | | | | |
| 58,305 | 53,290 | 24,565 | | | | | | | |
| 58,485 | 45,295 | 54,585 | | | | | | | |
| 0,490 | 0,300 | 54,625 | | | | | | | |
| | 0,560 | 40,655 | | | | | | | |
| | 48,565 | 0,660 | | | | | | | |
| | 0,580 | | | | | | | | |

| | | | | | | |
|---------|---------|----------|----------|----------|---------|-------|
| F0 | F1 | F2 | F3 | F4 | FNZ | AN |
| 0,5 | 450,5 | 1450,5 | 2450,5 | 3300,5 | 250,5 | 0,5 |
| 0,205 | 450,250 | 1450,250 | 2450,250 | 3300,250 | 250,830 | 0,830 |
| 120,295 | 200,285 | 1380,285 | 2470,285 | 3050,285 | | |
| 120,345 | 409,305 | 1400,305 | 2470,305 | 3050,305 | | |
| 95,415 | 620,350 | 850,365 | 2570,395 | 3300,355 | | |
| 60,475 | 620,435 | 850,420 | 2570,405 | | | |
| 0,485 | 461,485 | 1343,485 | 2018,485 | | | |
| 0,535 | 300,505 | 1720,505 | 2250,505 | | | |
| 55,540 | 300,565 | 1680,565 | 2250,570 | | | |
| 55,735 | 340,580 | 1680,590 | 2150,590 | | | |
| 0,740 | 230,640 | 1212,660 | 2258,660 | | | |
| | 203,645 | 1212,735 | 2258,735 | | | |
| | 203,735 | 1450,740 | 2450,740 | | | |
| | 450,740 | | | | | |
| A1 | A2 | A3 | A4 | A5 | A6 | AB |
| 0,5 | 0,5 | 0,5 | 0,5 | 0,5 | 0,5 | 0,5 |
| 0,830 | 0,560 | 0,260 | 0,260 | 0,260 | 0,260 | 0,830 |
| | 54,565 | 56,265 | 58,265 | 48,265 | 36,265 | |
| | 54,735 | 56,325 | 58,325 | 48,325 | 36,325 | |
| | 0,740 | 0,330 | 0,330 | 0,330 | 0,330 | |
| | | 0,560 | 0,560 | 0,560 | | |
| | | 30,565 | 43,565 | 46,565 | | |
| | | 30,735 | 43,735 | 46,735 | | |
| | | 0,740 | 0,740 | 0,740 | | |
| B1 | B2 | B3 | | | | |
| 50,0 | 70,5 | 110,0 | | | | |
| 50,250 | 70,250 | 110,250 | | | | |
| 70,275 | 120,275 | 180,275 | | | | |
| 70,415 | 120,305 | 180,305 | | | | |
| 300,445 | 50,375 | 140,355 | | | | |
| 300,735 | 50,415 | 140,415 | | | | |
| 50,740 | 250,445 | 300,455 | | | | |
| | 250,735 | 300,735 | | | | |
| | 0,735 | 0,740 | | | | |

Appendix B. Synthesizer parameters for the baseline stimuli were identical to the default output from the IPOX interface (Dirksen & Coleman, 1995) with the exception of /k_{ɒb}/ and /k_{ɒp}/.

For these stimuli, the following changes were made:

| CVC | Klatt Parameter | Time point (ms) | Value |
|--------------------|-----------------|-----------------|-------|
| /k _{ɒb} / | AB | ~650 | 0 |
| | A2 | 650-700 | 20 |
| | A3 | 650-700 | 30 |
| | F2 | 430 | 850 |
| | F2 | 830 | 700 |
| /k _{ɒp} / | AB | ~565 | 0 |
| | A2 | 565-640 | 50 |
| | A3 | 565-640 | 40 |

Effect of Time and Frequency Manipulation on Syllable Perception in Developmental Dyslexics

Ken I. McAnally, Peter C. Hansen, Piers L. Cornelissen, and John F. Stein
J Speech Lang Hear Res 1997;40:912-924

This article has been cited by 6 article(s) which you can access for free at:
<http://jslhr.asha.org/cgi/content/abstract/40/4/912#otherarticles>

This information is current as of April 12, 2010

This article, along with updated information and services, is
located on the World Wide Web at:
<http://jslhr.asha.org/cgi/content/abstract/40/4/912>



AMERICAN
SPEECH-LANGUAGE-
HEARING
ASSOCIATION